

MASTIRID®

Protect udder with Mastirid

Your best shoice to combat Mastitis







MASTIRID

A broad spectrum combination of Cephalexin & Neomycin with Prednisolone

Cephalexin a bactericidal antibiotic belonging to the Cephalosporin group. It is effective against most of the gram positive bacteria Viz - Staphylococcus aureus, Streptococcus agalactiae, including resistant varieties of penicillin (penicillinase positive) that are main causes of mastitis, and moderately effective against gram negative bacteria.

Neomycin is a aminoglycoside antibiotic and has an excellent activity against Gram negative bacteria Viz - E.Coli, Proteus spps., Klebsiella spps., Salmonella spps., etc. and has a moderate activity against Gram positive bacteria.

Prednisolone gives a quick local anti-inflammatory effect that allows a more effective reach of antibiotic to the infection site.

DOSAGE:

Cattle:

One syringe (5 gm) per affected quarter every 12 hrs for 3 times or as directed by veterinarian.

Sheep & Goat:

Half syringe (2.5 gm) per affected quarter every 12 hrs for 3 times or as directed by veterinarian.



Each 5 gm tube contains:

Cephalexin IP equivalent to
Cephalexin (anhydrous) 100 mg
Neomycin Sulphate IP 100 mg
Prednisolone IP 10 mg



MASTIRID